

## X-Ray Quiz: Ankle Sprain

Elif Kılıçlı, Meliha Fındık, Cemil Kavalcı

Department of Emergency Medicine, Başkent University Faculty of Medicine, Ankara, Turkey

### Introduction

A 55-year-old man presented to the our emergency department (ED) with a complaint of ankle pain after a foot sprain. Initial evaluations revealed a swollen foot, tenderness, and limited active range of motion of the ankle. Standard posteroanterior (PA) and lateral X-rays of the ankle were performed (Figure 1-2).

### Questions

1. What are the radiological abnormalities seen in Figure 1-2?
2. What is the diagnosis?
3. How should this case be managed?



**Figure 1.** Dislocation of lisfranc joint and fracture of the third metatars basis on X-ray



**Figure 2.** Dislocation of lisfranc joint and fracture of the third metatars basis on X-ray

Answer: p. 44

**Correspondence to:** Elif Kılıçlı, Department of Emergency Medicine, Başkent University Faculty of Medicine, Ankara, Turkey  
Phone: +90 312 212 59 83 e.mail: kiliclie@mynet.com

**Received:** 04.02.2014 **Accepted:** 20.02.2014

©Copyright 2014 by Emergency Physicians Association of Turkey - Available online at [www.akademikaciltip.com](http://www.akademikaciltip.com)  
DOI:10.5152/jaem.2014.152

