

Development of Pituitary Apoplexy in a Patient with Meningioma and Pituitary Macroadenoma: A Case Report

© Hakan Yılmaz¹, © Zelal Adıbelli², © Canan Akkuş², © Hakan Demirci², © Bülent Mızrak³, © Cevdet Duran⁴

¹Uşak University Faculty of Medicine, Department of Radiology, Uşak, Turkey

²Uşak University Faculty of Medicine, Department of Internal Medicine, Uşak, Turkey

³Uşak University Faculty of Medicine, Department of Pathology, Uşak, Turkey

⁴Uşak University Faculty of Medicine, Department of Internal Medicine, Division of Endocrinology and Metabolism, Uşak, Turkey

Abstract

Although pituitary adenomas and meningiomas are among the commonly encountered benign tumors, the co-existence of these tumors is rare. One of the most important complications of pituitary adenomas is the development of apoplexy, often resulting in death if left untreated. In this article, we present a patient admitted to the emergency department with complaints of sudden-onset headache, nausea, vomiting, and optic nerve paralysis. On examination, the case was found to have parietal lobe meningioma and pituitary adenoma. Therefore, the case underwent emergent transsphenoidal surgery considering pituitary apoplexy and was treated with l-thyroxine and hydrocortisone in the post-operative period. Although the co-existence of a pituitary adenoma and a meningioma is known in the literature, such a co-existence with apoplexy is the first case to be described in the literature.

Keywords: Apoplexy, hypophysis, meningioma, pituitary adenoma

Introduction

Pituitary apoplexy is one of the major complications of pituitary adenomas and is known to be one of the endocrinological emergencies manifesting itself with headache, visual impairment, hypopituitarism, and cranial nerve paralysis. If untreated medically and/or surgically, pituitary apoplexy can result in death. The frequency of apoplexy ranges between 2 and 12% in patients with pituitary adenomas (1). Apoplexy generally develops suddenly at the base of a non-functional or functional pituitary adenoma, triggered by hypertension, anticoagulant therapy, increased intracranial pressure, and major surgery, and sometimes also occurs idiopathically due to necrosis triggered by dynamic tests and sometimes owing to hemorrhages, and thus resulting in a rapid increase in the size of the pituitary gland (1,2). Apoplexy may also lead to deaths if it goes untreated surgically and/or conservatively medically.

Meningiomas are tumors with a commonly often benign nature and originating from arachnoidal cap cells by constituting 13%-26% of intracranial tumors (3). Meningiomas are generally seen in later periods of life and among women (3). Complete surgical excision is the standard treatment, and radiotherapy can also be administered for treating atypical, recurrent, or malignant meningiomas (3).

Although the number of meningiomas is limited, the cases of meningiomas co-existent with functional or non-functional pituitary adenomas have been reported in the literature (4-6). Some cases of meningiomas, especially those originating from the region of the diaphragmatic sella, may present with an appearance similar to pituitary adenomas and cause diagnostic confusion (4-6).

In our report, we presented a case detected with a pituitary macroadenoma and a right parietal lobe meningioma in



Corresponding Author: Cevdet Duran MD, Uşak University Faculty of Medicine, Department of Internal Medicine, Division of Endocrinology and Metabolism, Uşak, Turkey
Phone: +90 533 429 85 85 **E-mail:** drcduran@gmail.com **ORCID ID:** orcid.org/0000-0001-9560-180X

Cite this article as: Yılmaz H, Adıbelli Z, Akkuş C, Demirci H, Mızrak B, Duran C. Development of Pituitary Apoplexy in a Patient with Meningioma and Pituitary Macroadenoma: A Case Report. Eurasian J Emerg Med. 2023;22(1): 58-61.

©Copyright 2023 by the Emergency Medicine Physicians' Association of Turkey
Eurasian Journal of Emergency Medicine published by Galenos Publishing House.

Received: 19.06.2021
Accepted: 06.08.2021

emergency examinations performed due to the clinic of pituitary apoplexy, where hypopituitarism developed after surgical debulking by the transsphenoidal route.

Case Report

A 72-year-old female patient was found to develop severe headache with sudden onset, nausea, vomiting, vision loss in the left eye, and left-eyelid ptosis on cranial computerized tomography (CT) performed nearly three months ago (on 8th October 2020). On CT examination, an extra-axially located calcific meningioma on the right side of the cranium was determined (Figures 1A, B). Additionally, a peripherally located extra-axial meningioma in size of 12x11 mm was detected in the lateral part of the parietal lobe on cranial magnetic resonance imaging (MRI). A 22 mm adenoma, the borders of which could not be distinguished, was also identified within the pituitary gland (Figures 2A-D). At once, considering pituitary adenoma and apoplexy in the case, an emergent pituitary decompression surgery was performed by the transsphenoidal route under the coverage of steroids, and during the surgery, the pressurized hematoma or adenoma was observed to be evacuated after the dura incision. The histopathological examination of the hypophysectomy material of the case, where postoperative hypopituitarism was detected and steroid therapy was continued, revealed that the lesion was completely necrotic. Through the medical history, it was determined that the case had been diagnosed with low-grade non-Hodgkin lymphoma (NHL) 34 years ago, received chemotherapy due to the recurrence twice 34 and 12 years ago, undergone hysterectomy 30 years ago, had hypothyroidism for 25 years, and so received the treatment with levothyroxine 50 µg/day po recently, and had chronic hepatitis B infection for 18 years and thus been treated with entecavir of 0.5 mg tb 1x1 po lately. The medical history also revealed that the case had hypertension for 10 years, and most recently received a combination therapy of perindopril/indapamide 4 mg/1.25 mg 1x1 po with amlodipine of 5 mg 1x1 po, as well as receiving metformin 2x850 mg po due to type 2 diabetes mellitus




Figure 1. The extra-axially located calcific nodular mass lesion (calcific meningioma) is present on the right side of the bone window (A) and parenchymal window images (B) taken through unenhanced computerized tomography

(T2DM) for the last seven years. The patient having received hydrocortisone treatment of 3x10 mg po within the latest period was admitted to our department. On physical examination, the case (height: 162 cm, weight: 72 kg, and body mass index: 27.4 kg/m²), was found to be conscious, cooperative, and full of motor functions. Blood pressure and visual field examination of the case were observed to be within the normal limits. Given the re-examination of the specimens prepared in the pathology unit, it was detected that while the preparations were inappropriate for healthy histopathological evaluation due to necrosis, the background was composed of the cells with monotonous appearance not constituting a distinct pattern, and these cells were stained with synaptophysin in immunohistochemical examination (Figures 3A, B). Therefore, the case was followed up in the outpatient clinic with the current treatment.

Discussion

In this report, we present a case developing pituitary apoplexy at the base of a pituitary adenoma and a simultaneous meningioma detected in the right parietal region. To our knowledge, no reports, including the co-existence of pituitary apoplexy and a meningioma, were encountered in the literature. Albeit their rarity, pituitary adenomas and meningiomas can be seen together. In some cases, both conditions can be found together as a collision tumor (7). Since adenomas and meningiomas are




Figure 2. A) T2-weighted sagittal section. Magnetic resonance imaging (MRI) shows the heterogeneous enlargement of the pituitary gland, (B) T1-weighted contrast-enhanced sagittal, and (C) T1-weighted non-contrast coronal section. MRI shows the enlargement of the pituitary gland with heterogeneous hyperintense hemorrhagic areas, and (D) T1-weighted contrast-enhanced coronal section. MRI shows the heterogeneous contrast enhancement of the enlarged pituitary gland




Figure 3. A) The area containing necrobiotic cells surrounded by necrotic tissue and thrombus-including venule (x20 magnification, stained with hematoxylin and eosin), and (B) diffuse synaptophysin positivity in cells. The vessel wall was negatively stained (x40 magnification)

often required to be removed by transsphenoidal surgery and craniotomy, respectively, the preoperative differential diagnoses of pituitary adenomas and meningiomas, particularly arising from diaphragmatic or tuberculum sella, are so important (6). It has been reported that contrast-enhanced dynamic pituitary MRI can be beneficial in the differential diagnosis of these two conditions (5). The meningioma in our case was detected in the parietal region, and thus, there was no diagnostic confusion. The reason for the co-existence of a meningioma and a pituitary adenoma has yet to be fully elucidated. In a study comparing 57 cases with pituitary adenoma-related meningiomas with those sporadic pituitary adenomas and sporadic meningiomas, Zhu et al. (8) reported that the lower expression of the multiple endocrine neoplasia type 1 gene is central to pituitary adenoma-related meningiomas by upregulating the mammalian target of the rapamycin (mTOR) signaling pathway, and that the treatment with rapamycin, an mTOR inhibitor, can be used by increasing apoptosis in treating pituitary adenomas in the future.

In our case, such clinically pre-existing symptoms as headache, problems of vision, and galactorrhea were absent. Additionally, our case was deprived of other clinical findings suggesting Cushing's syndrome or acromegaly, and probably the adenoma in our case was also non-functional. On pathological examination, it was determined that the base was composed of cells with a monotonous appearance not forming a distinct pattern. The immunohistochemical examination also revealed that these cells were stained with synaptophysin. Even so, no immunostaining could be performed for pituitary hormones due to necrotic tissues. The presence of necrotic tissues in the surgical material was also compatible with apoplexy. In her history, our case was determined to have been diagnosed with NHL, as in a remission state. Base on the literature, NHL exhibits sometimes an adenoma-like image by leading to the pituitary

infiltration (9); however, there was no finding suggesting the lymphoma infiltration in the pathological examination of our case.

The prevalence of apoplexy ranges between 2-12% in patients with pituitary adenomas (1). Apoplexy is an emergency-requiring condition developing usually with hypertension, anticoagulant therapy, etc. at the base of a non-functional or functional pituitary adenoma, or developing sometimes idiopathically, occurring suddenly after the necrosis or hemorrhages, and resulting in a rapid increase in the size of the pituitary gland; it may also result in deaths if untreated surgically and/or conservatively medically (1,2). Her history revealed that our case had a medical history of hypertension and T2DM, and so receiving three antihypertensive drugs and one oral antidiabetic medication per day due to these diseases. The case also received L-thyroxine therapy for the pre-existing hypothyroidism. When diagnosed with apoplexy, the case underwent transsphenoidal surgery under the coverage of steroids and as the continuation therapy, hydrocortisone combined with L-thyroxine was continued.

Our case had a history of NHL existing for many years and currently in remission. Based on the literature, various cases rarely developing apoplexy in the lymphoma background have been reported (10). However, in our case, there was no lymphoma infiltration in the postoperative tissue, and immunostaining with synaptophysin was consistent with the pituitary adenoma.

Conclusion

In conclusion, in the report, we presented a case developing the pituitary apoplexy at the base of a pituitary adenoma and a parietal meningioma. Although the co-existence of a pituitary adenoma and a meningioma is known in the literature, such a co-existence with apoplexy is the first case to be described in the literature.

Ethics

Informed Consent: Consent form was filled out by all participants.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices - Concept - Design - Data Collection or Processing - Analysis or Interpretation - Literature Search - Writing: H.Y., Z.A., C.A., H.D., B.M., C.D.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Glezer A, Bronstein MD. Pituitary apoplexy: pathophysiology, diagnosis and management. *Arch Endocrinol Metab.* 2015;59:259-64.
2. Briet C, Salenave S, Bonneville JF, Laws ER, Chanson P. Pituitary Apoplexy. *Endocr Rev.* 2015;36:622-45.
3. Marosi C, Hassler M, Roessler K, Reni M, Sant M, Mazza E, et al. Meningioma. *Crit Rev Oncol Hematol.* 2008;67:153-71.
4. Amirjamshidi A, Mortazavi SA, Shirani M, Saeedinia S, Hanif H. 'Coexisting pituitary adenoma and suprasellar meningioma-a coincidence or causation effect: report of two cases and review of the literature'. *J Surg Case Rep.* 2017;2017:rjx039.
5. Cappabianca P, Cirillo S, Alfieri A, D'Amico A, Maiuri F, Mariniello G, et al. Pituitary macroadenoma and diaphragma sellae meningioma: differential diagnosis on MRI. *Neuroradiology.* 1999;41:22-6.
6. Taylor SL, Barakos JA, Harsh GR 4th, Wilson CB. Magnetic resonance imaging of tuberculum sellae meningiomas: preventing preoperative misdiagnosis as pituitary macroadenoma. *Neurosurgery.* 1992;31:621-7.
7. Zhao Y, Zhang H, Lian W, Xing B, Feng M, Liu X, et al. Collision tumors composed of meningioma and growth hormone-secreting pituitary adenoma in the sellar region: Case reports and a literature review. *Medicine (Baltimore).* 2017;96:e9139.
8. Zhu H, Miao Y, Shen Y, Guo J, Xie W, Zhao S, et al. The clinical characteristics and molecular mechanism of pituitary adenoma associated with meningioma. *J Transl Med.* 2019;17:354.
9. Tamer G, Kartal I, Aral F. Pituitary infiltration by non-Hodgkin's lymphoma: a case report. *J Med Case Rep.* 2009;3:9293.
10. Rizek P, Seitelbach M, Alturkustani M, Leung A, Fraser JA. Sellar and parasellar intravascular lymphoma mimicking pituitary apoplexy. *J Neuroophthalmol.* 2012;32:33-7.